Communications in Contemporary Mathematics
(2015) 1550062 (18 pages)
© World Scientific Publishing Company
DOI: 10.1142/S0219199715500625

World Scientific
www.worldscientific.com

On the spin-1 Bose–Einstein condensates in the presence of Ioffe–Pritchard magnetic field

Hichem Hajaiej
Department of Mathematics, College of Sciences
King Saud University, 11451 Riyadh, Saudi Arabia
hhajaiej@ksu.edu.sa

Rémi Carles
CNRS et Université Montpellier, Mathématiques
CC 051, Place Eugène Bataillon
34095, Montpellier Cedex 5, France
remi.carles@math.cnrs.fr

Received 15 August 2014
Revised 22 April 2015
Accepted 8 June 2015
Published 7 July 2015

We study the Cauchy problem of an antiferromagnetic spin-1 Bose–Einstein condensate under Ioffe–Pritchard magnetic field B. We then address the existence of ground state solutions and characterize the orbit of standing waves.

Keywords: Nonlinear Schrödinger equation; magnetic field; Bose–Einstein condensation; ground state; standing waves.

Mathematics Subject Classification 2010: 35Q55, 35J20, 35B35

Commun. Contemp. Math. Downloaded from www.worldscientific.com
by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.

1. Introduction

In 1995, it was proved that massive noninteracting bosons at low temperature can occupy the same lowest energy (ground state solution) single particle state and form the Bose–Einstein condensate. The total spin number F corresponding to the lowest energy has to be an integer with $2F+1$ hyperfine states ($m_F = -F, -F+1, \ldots, F$). They are called spin-F (BEC).

The recent developments of optical trapping techniques enabled to confine atoms independently of their spin orientation to obtain the spinor condensates. The main motivation of such condensates lies essentially in their numerous applications in quantum mechanics. This cannot be covered by the classical single (BEC), which do not possess the appropriate properties.

In this paper, we will focus our study on the 1-spin (BEC) which is described by the following Gross–Pitaevskii system:

$$\begin{cases} i\partial_t\psi_1 = (H+\beta_s(|\psi_1|^2+|\psi_0|^2-|\psi_{-1}|^2))\psi_1+\beta_s\psi_0^2\bar{\psi}_{-1}+B\psi_0, \\ i\partial_t\psi_0 = (H+\beta_s(|\psi_1|^2+|\psi_{-1}|^2))\psi_0+2\beta_s\psi_1\bar{\psi}_0\psi_{-1}+\bar{B}\psi_1+B\psi_{-1}, \\ i\partial_t\psi_{-1} = (H+\beta_s(|\psi_{-1}|^2+|\psi_0|^2-|\psi_1|^2))\psi_{-1}+\beta_s\psi_0^2\bar{\psi}_1+\bar{B}\psi_0, \\ \psi_j(0,x) = \Phi_j(x) \quad \text{for any } j\in\{-1,0,1\}, \end{cases} \tag{1.1}$$

where $x\in\mathbb{R}^d, 1\le d\le 3$, and the (nonlinear) operator H is defined by

$$H = -\frac{1}{2}\Delta + V_{\text{ext}}(x) + \beta_n\sum_{j=-1}^{1}|\psi_j|^2.$$

The external potential is typically a harmonic potential, possibly anisotropic,

$$V_{\text{ext}}(x) = \sum_{j=1}^{d}\omega_j^2\frac{x_j^2}{2}. \tag{1.2}$$

The function $\psi_j(t,x)$ is the complex-valued wave function of the jth ($j=-1,0,1$) component of $\psi=(\psi_{-1},\psi_0,\psi_1)$. The parameters β_n and β_s describe the spin-independent interaction and spin-exchange interaction, respectively. They can be positive or negative. They are proportional to the total number of atoms N. They describe the collisions between atoms (see [1] and references therein). The function $B\in L^\infty(\mathbb{R}^d;\mathbb{C})$ denotes the external Ioffe–Pritchard magnetic field. For a more detailed account of the physical model, the reader can refer to [9, 11, 13].

As noticed in [1], the following quantities are formally conserved by the flow:

$$\text{Total mass: } \frac{d}{dt}\left(\sum_{j=-1}^{1}\|\psi_j(t)\|_{L^2}^2\right)=0, \tag{1.3}$$

$$\text{Total energy: } \frac{d}{dt}\left(\widehat{E}_0(\psi(t))+2\,\text{Re}\left(\int_{\mathbb{R}^d}B(\bar{\psi}_1\psi_0+\bar{\psi}_0\psi_{-1})dx\right)\right)=0, \tag{1.4}$$

where the magnetic-free energy is defined by

$$\widehat{E}_0(\psi)=\int_{\mathbb{R}^d}\left(\sum_{j=-1}^{1}\left(\frac{1}{2}|\nabla\psi_j|^2+V_{\text{ext}}(x)|\psi_j|^2\right)+\frac{\beta_n}{2}\left(\sum_{j=-1}^{1}|\psi_j|^2\right)^2\right.$$

$$\left.+\frac{\beta_s}{2}(|\psi_1|^2-|\psi_{-1}|^2)^2+\beta_s|\psi_0|^2(|\psi_1|^2+|\psi_{-1}|^2)+2\beta_s\,\text{Re}(\bar{\psi}_1\psi_0^2\bar{\psi}_{-1})\right)dx.$$

We denote by $\widehat{E}(\psi)$ the total energy, which is formally time-independent,

$$\widehat{E}(\psi)=\widehat{E}_0(\psi(t))+2\,\text{Re}\left(\int_{\mathbb{R}^d}B(\bar{\psi}_1\psi_0+\bar{\psi}_0\psi_{-1})dx\right). \tag{1.5}$$

Commun. Contemp. Math. Downloaded from www.worldscientific.com by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.

In the present study, we will consider the case where the spin interaction is repulsive, $\beta_n > 0$, and the spin exchange interaction is antiferromagnetic $\beta_s > 0$. The only observable solutions of (1.1) in experiments are the so-called ground state solutions. They are the minimizers of the following constrained variational problem.

For two prescribed positive numbers $M, N > 0$, we set

$$\widehat{I}_{M,N} = \inf_{\psi=(\psi_{-1},\psi_0,\psi_1)\in\widehat{C}_{M,N}} \widehat{E}(\psi), \tag{1.6}$$

where

$$\widehat{C}_{M,N} = \Bigg\{\psi = (\psi_{-1},\psi_0,\psi_1) \in \Sigma_{\mathbb{C}}(\mathbb{R}^d) \times \Sigma_{\mathbb{C}}(\mathbb{R}^d) \times \Sigma_{\mathbb{C}}(\mathbb{R}^d):$$

$$\int_{\mathbb{R}^d} (|\psi_{-1}|^2 + |\psi_0|^2 + |\psi_1|^2)dx = N \quad \text{and}$$

$$\int_{\mathbb{R}^d} (|\psi_1|^2 - |\psi_{-1}|^2)dx = M\Bigg\}, \tag{1.7}$$

with

$$\Sigma(\mathbb{R}^d) = \Sigma = \left\{u \in H^1(\mathbb{R}^d;\mathbb{R}) : |u|^2_{V_{\text{ext}}} := \int V_{\text{ext}}(x)u(x)^2dx < \infty\right\}$$

and

$$|u|^2_{\Sigma} = \|u\|^2_{L^2} + \|\nabla u\|^2_{L^2} + |u|^2_{V_{\text{ext}}}.$$

$\Sigma_{\mathbb{C}}$ denotes the set of complex-valued functions with a Σ regularity:

$$\Sigma_{\mathbb{C}}(\mathbb{R}^d) = \widehat{\Sigma} = \{z = (u,v) \simeq u + iv; (u,v) \in \Sigma \times \Sigma\}.$$

In terms of the polar coordinates $\psi_j = |\psi_j|e^{i\theta_j}$, $\hat{E}_0$ can be rewritten in the following way:

$$\widehat{E}_0(\psi) = \int_{\mathbb{R}^d} \Bigg(\sum_{j=-1}^{1} \left(\frac{1}{2}|\nabla\psi_j|^2 + V_{\text{ext}}(x)|\psi_j|^2\right) + \frac{\beta_n}{2}\left(\sum_{j=-1}^{1}|\psi_j|^2\right)^2$$

$$+ \frac{\beta_s}{2}(|\psi_1|^2 - |\psi_{-1}|^2)^2 + \beta_s|\psi_0|^2(|\psi_1|^2 + |\psi_{-1}|^2)$$

$$+ 2\beta_s|\psi_1||\psi_0|^2|\psi_{-1}|\cos(2\theta_0 - \theta_1 - \theta_{-1})\Bigg)dx. \tag{1.8}$$

Let θ_B be the angle function such that $B = |B|e^{i\theta_B}$.

$$\widehat{E}(\psi) = \int_{\mathbb{R}^d} \Bigg(\sum_{j=-1}^{1} \left(\frac{1}{2}|\nabla\psi_j|^2 + V_{\text{ext}}(x)|\psi_j|^2\right) + \frac{\beta_n}{2}\left(\sum_{j=-1}^{1}|\psi_j|^2\right)^2$$

$$+ \frac{\beta_s}{2}(|\psi_1|^2 - |\psi_{-1}|^2)^2 + \beta_s|\psi_0|^2(|\psi_1|^2 + |\psi_{-1}|^2)$$

Commun. Contemp. Math. Downloaded from www.worldscientific.com by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.

$$+2\beta_s|\psi_1||\psi_0|^2|\psi_{-1}|\cos(2\theta_0-\theta_1-\theta_{-1})$$

$$+2|B||\psi_0|(|\psi_1|\cos(\theta_B+\theta_0-\theta_1)+|\psi_{-1}|\cos(\theta_B+\theta_{-1}-\theta_0))\Bigg)dx. \tag{1.9}$$

To (1.9) (i.e. (1.5)), we will associate the following constrained minimization problem

$$I_{M,N}=\inf_{u=(u_{-1},u_0,u_1)\in C_{M,N}} E(u), \tag{1.10}$$

where

$$C_{M,N}=\Bigg\{u=(u_{-1},u_0,u_1)\in\Sigma^3:\int_{\mathbb{R}^d}(u_{-1}^2+u_0^2+u_1^2)=N \text{ and}$$

$$\int_{\mathbb{R}^d}(u_1^2-u_{-1}^2)=M\Bigg\},$$

$$E_0(u)=\frac{1}{2}\|\nabla u\|_2^2+\int_{\mathbb{R}^d}\Bigg(\sum_{j=-1}^{1}V_{\text{ext}}(x)u_j^2+\frac{\beta_n}{2}\Bigg(\sum_{j=-1}^{1}u_j^2\Bigg)^2 \tag{1.11}$$

$$+\frac{\beta_s}{2}(u_1^2-u_{-1}^2)^2+\beta_s u_0^2(u_1^2+u_{-1}^2)-2\beta_s u_1u_0^2u_{-1}\Bigg)dx$$

and

$$E(u)=E_0(u)-2\int_{\mathbb{R}^d}|B(x)||u_0||u_1|-2\int_{\mathbb{R}^d}|B(x)||u_0||u_{-1}|. \tag{1.12}$$

Note that (1.11) can be rewritten in the following way:

$$E_0(u)=\frac{1}{2}\|\nabla u\|_2^2+\int_{\mathbb{R}^d}\sum_{j=-1}^{1}V_{\text{ext}}(x)u_j^2(x)+\frac{\beta_n+\beta_s}{2}\int_{\mathbb{R}^d}(u_1^4+u_{-1}^4)dx$$

$$+\frac{\beta_n}{2}\int_{\mathbb{R}^d}u_0^4+(\beta_n+\beta_s)\int_{\mathbb{R}^d}u_0^2(u_1^2+u_{-1}^2)dx$$

$$+(\beta_n-\beta_s)\int_{\mathbb{R}^d}u_1^2u_{-1}^2dx-2\beta_s\int u_1u_{-1}u_0^2dx. \tag{1.13}$$

Recall that (1.1) has been studied numerically quite recently [1]. The authors have been able to show the existence of ground state solutions of (1.5) via some readapted and innovative numerical methods. Stationary states with lowest energy are very important in physics, since they enable us to understand the properties of BEC. In the literature, there exist many results, based on experiments, proving the existence of ground states of 1-spin condensates, see, e.g., [1, 3, 14] and references therein.

Commun. Contemp. Math. Downloaded from www.worldscientific.com by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.

To our knowledge, the first theoretical result established in this context, has been established in [3]. However the authors only considered the case $d = 1, V_{\text{ext}} \equiv 0$ and $B \equiv 0$. This significantly limits the scope of applications in physics. They proved the existence of ground state solutions when $\beta_n < 0$ and $\beta_s < 0$. When $V_{\text{ext}} \neq 0$, we will prove that (1.10) has a minimizer when both β_n and β_s are non-negative. Note that in this case, both spin-independent interaction and spin-exchange interaction are repulsive, and if there is no trapping potential V_{ext}, the atoms cannot be confined, this is why when β_n and $\beta_s > 0$ and $V_{\text{ext}} \equiv 0$, there are no ground state solutions. Technically, (1.10) is a problem which is completely different in the presence or in the absence of the trapping potential V_{ext}.

When $V_{\text{ext}} \in L^{\infty}_{\text{loc}}(\mathbb{R}^d)$ and $\lim_{R\to\infty}(\operatorname{ess\,inf}_{|x|\geq R} V(x)) = \infty$, Lin and Chern [14] have studied (1.6) with $B \equiv 0$. They were essentially concerned with the bifurcation problem of the ground states. More precisely, they have studied the bifurcation problem:

$$\begin{cases} (\mu+\lambda)u_1 = Hu_1 + 2\beta_s(u_0^2(u_1-u_{-1}) + u_1(u_1^2-u_{-1}^2)), \\ \mu u_0 = Hu_0 + 2\beta_s u_0(u_1-u_{-1})^2, \\ (\mu-\lambda)u_1 = Hu_1 + 2\beta_s(u_0^2(u_{-1}-u_1) + u_{-1}(u_{-1}^2-u_1^2)), \end{cases}$$

where λ and μ are the Lagrange multipliers arising from the constraint $C_{M,N}$.

In this paper, we give a complete study of the spin-1 (BEC) problem. First, we prove in Sec. 3 that the Cauchy problem (1.1) has a unique solution in a suitable function space. In Sec. 4, we characterize the orbit of ground state solutions. To achieve this goal, we will need the results established in Sec. 2, where we prove the existence of these stationary solutions. Our method hinges on the approach built up by Cazenave and Lions in [8], and by Hajaiej–Stuart in [10]. In [8], the authors gave a precise line of attack to establish the orbital stability. In fact, they showed that it is sufficient to show that all minimizing sequences of the variational problem (1.6) are relatively compact in $\hat{\Sigma}^3$. To characterize the orbit of standing waves, a more subtle and in-depth study must be undertaken. This has been developed in [10]. In Sec. 4, we give a detailed account for the current framework.

2. Notations

For fixed positive numbers M and N, we define

$$Z_{M,N} = \{z = (z_{-1}, z_0, z_1) \in \widehat{C}_{M,N} : \widehat{E}(z) = \widehat{I}_{M,N}\} \tag{2.1}$$

and

$$W_{M,N} = \{u = (u_{-1}, u_0, u_1) \in C_{M,N} : u_{-1}, u_0 \text{ and } u_1 > 0 \text{ and } E(u) = I_{M,N}\}.$$

Definition 2.1. We say that $Z_{M,N}$ is stable if $Z_{M,N} \neq \emptyset$, and for all $w \in Z_{M,N}$, for any $\varepsilon > 0$, there exists $\delta > 0$ such that for any $\psi^0 \in \widehat{\Sigma}^3$ satisfying $\|\psi^0 - w\|_{\widehat{\Sigma}^3} < \delta$,

Commun. Contemp. Math. Downloaded from www.worldscientific.com by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.

then for all $t \in \mathbb{R}$,

$$\inf_{z \in Z_{M,N}} \|\psi(t,\cdot) - z\|_{\widehat{\Sigma}^3} < \varepsilon, \tag{2.2}$$

where $\psi(t,\cdot)$ is the unique solution of (1.1) initial condition ψ^0.

The following result can be found in, e.g., [12].

Lemma 2.2. *Σ is compactly embedded in $L^q(\mathbb{R}^d)$ for any $2 \le q < \frac{2d}{d-2}$.*

Lemma 2.3. (1) *The minimization problem* (1.10) *is well-posed and any minimizing sequence of* (1.10) *is bounded in Σ^3.*

(2) *Any minimizing sequence of* (1.10) *is relatively compact in Σ^3, i.e. for all sequences $u_n \in C_{M,N}$ such that $E(u_n) \to I_{M,N}$, there exists $u \in \Sigma^3$ such that, up to a subsequence, $u_n \to u$ in Σ^3.*

(3) *The functionals E and $\widehat{E}$ are C^1 on Σ^3 and $\widehat{\Sigma}^3$, respectively.*

(4) *The function:*

$$(0,\infty) \times (0,\infty) \to \mathbb{R}$$
$$(M,N) \mapsto I_{M,N}$$

is continuous with respect to both variables.

Proof. (1) Since we are supposing that β_n and $\beta_s \ge 0$, we can easily deduce that

$$\begin{aligned} E(u) &\ge -2\int_{\mathbb{R}^d} |B(x)||u_0||u_1| + |B(x)||u_0||u_{-1}|dx \\ &\ge -2\|B\|_\infty \|u_0\|_{L^2} \left(\|u_1\|_{L^2} + \|u_{-1}\|_{L^2}\right) \\ &\ge -\|B\|_{L^\infty} \left(\|u_0\|_{L^2}^2 + \|u_1\|_{L^2}^2 + \|u_0\|_{L^2}^2 + \|u_{-1}\|_{L^2}^2\right) \ge -2\|B\|_\infty N. \end{aligned}$$

This proves that the minimization problem $I_{M,N}$ is well-posed and that every minimizing sequence is bounded in Σ^3.

(2) Let $u_n = (u_{n,-1}, u_{n,0}, u_{n,1})$ be a minimizing sequence of (1.10). From (1), we know that (u_n) is bounded in Σ^3, therefore (up to a subsequence) there exists $u = (u_{-1}, u_0, u_1) \in \Sigma^3$ such that $u_{n,j} \rightharpoonup u_j$ in $\Sigma, -1 \le j \le 1$.

First note that by the lower semi-continuity of the norm Σ we have

$$|u|_{\Sigma^3}^2 \le \liminf |u_n|_{\Sigma^3}^2. \tag{2.3}$$

By Lemma 2.2, we have $u_{n,j} \to u_j$ in $L^p(\mathbb{R}^d)$ for any $2 \le p < \frac{2d}{d-2}$ and $-1 \le j \le 1$.

In particular with $p = 4$, we infer

$$\lim_{n\to\infty} \int_{\mathbb{R}^d} (u_{n,1}^2 - u_{n,-1}^2)^2 = \int_{\mathbb{R}^d} (u_1^2 - u_{-1}^2)^2, \tag{2.4}$$

Commun. Contemp. Math. Downloaded from www.worldscientific.com by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.

$$\lim_{n\to+\infty}\int_{\mathbb{R}^d} u_{n,0}^2(u_{n,1}^2+u_{n,-1}^2)=\int_{\mathbb{R}^d} u_0^2(u_1^2+u_{-1}^2), \tag{2.5}$$

$$\lim_{n\to\infty}\int_{\mathbb{R}^d} u_{n,0}^2 u_{n,-1}u_{n,1}=\int_{\mathbb{R}^d} u_1u_0^2u_{-1}. \tag{2.6}$$

On the other hand, using the fact that $B\in L^\infty(\mathbb{R}^d)$, and that $u_{n,j}\to u_j$ in $L^2(\mathbb{R}^d)$, it follows that

$$\begin{aligned}\lim_{n\to+\infty}\int_{\mathbb{R}^d}|B|(|u_{n,0}||u_{n,1}|+|u_{n,0}||u_{n,-1}|)dx\\ =\int_{\mathbb{R}^d}|B|(|u_0||u_1|+|u_0||u_{-1}|)dx.\end{aligned} \tag{2.7}$$

Combining (2.3)–(2.7), we certainly have that

$$E(u)\le\liminf_{n\to\infty}E(u_n)=I_{M,N}.$$

But since $u=(u_{-1},u_0,u_1)\in C_{M,N}$, we can conclude that

$$E(u)=I_{M,N}.$$

(3) The proof is identical to [10, Proposition 3.2.9].
(4) The proof also goes exactly in the same way as in [10, Proposition 3.2.9]. Note also that in [14, Sec. 3] the continuity and monotonicity of $I_{M,N}$ have been discussed in details. □

Remark 2.4. Contrary to the case $V_{\text{ext}}\equiv 0$ studied in [3], it seems that it is not reasonable to expect to have a symmetric minimizer of (1.10), even when V_{ext} is isotropic. This is essentially due to the conflicts between the different terms of the energy functional. More precisely, note that $E(|u|)\le E(u)$, therefore without loss of generality, we can suppose that the minimizer of (1.10) is non-negative. Let u^* be its Schwarz symmetrization. Then using rearrangement inequalities established in [2], we certainly have that

$$\|\nabla u_j^*\|_{L^2}\le\|\nabla u_j\|_{L^2}\quad\text{for all } -1\le j\le 1, \tag{2.8}$$

$$\int_{\mathbb{R}^d}V_{\text{ext}}(x)(u_j^*)^2dx\le\int_{\mathbb{R}^d}V_{\text{ext}}(x)(u_j)^2dx\quad\text{for all } -1\le j\le 1, \tag{2.9}$$

$$(\beta_n+\beta_s)\int_{\mathbb{R}^d}(u_1^4+u_{-1}^4)dx=(\beta_n+\beta_s)\int_{\mathbb{R}^d}((u_1^*)^4+(u_{-1}^*)^4)dx, \tag{2.10}$$

$$\beta_n\int_{\mathbb{R}^d}u_0^4dx=\beta_n\int_{\mathbb{R}^d}(u_0^*)^4dx, \tag{2.11}$$

$$(\beta_n+\beta_s)\int_{\mathbb{R}^d}u_0^2(u_1^2+u_{-1}^2)dx\le(\beta_n+\beta_s)\int_{\mathbb{R}^d}(u_0^*)^2((u_{-1}^*)^2+(u_1^*)^2)dx. \tag{2.12}$$

Commun. Contemp. Math. Downloaded from www.worldscientific.com by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.

If in addition, $\beta_n < \beta_s$, then

$$(\beta_n - \beta_s) \int_{\mathbb{R}^d} u_1^2 u_{-1}^2 \geq (\beta_n - \beta_s) \int_{\mathbb{R}^d} (u_1^*)^2 (u_{-1}^*)^2, \tag{2.13}$$

$$\beta_s \int u_1 u_{-1} u_0^2 \leq \beta_s \int u_1^* u_{-1}^* (u_0^*)^2. \tag{2.14}$$

A classical tool to prove that (1.10) has a Schwarz symmetric minimizer is to show that $E(u^*) \leq E(u)$. Here (2.8), (2.9) and (2.13) are in conflict with (2.12) and (2.14). We have strong reasons to believe that there is no Schwarz symmetric minimizer for (1.10). This issue will be investigated.

From now on in an integral where no domain of integration is indicated, it is to be understood that it runs over $\mathbb{R}^d$. Sometimes we will omit the measure element dx.

Commun. Contemp. Math. Downloaded from www.worldscientific.com by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.

3. Cauchy Problem

Denote by

$$H_0 = -\frac{1}{2}\Delta + V_{\text{ext}}(x)$$

the linear part of H.

Definition 3.1. A pair (p, q) is admissible if $2 \leq q < \frac{2d}{d-2}$ ($2 \leq q \leq \infty$ if $d = 1, 2$ $\leq q < \infty$ if $d = 2$) and

$$\frac{2}{p} = \delta(q) := d\left(\frac{1}{2} - \frac{1}{q}\right).$$

We have the standard result (see, e.g., [7]).

Lemma 3.2. *Let $d \geq 1$, and suppose that the external potential is smooth and at most quadratic*:

$$\partial^\alpha V_{\text{ext}} \in L^\infty(\mathbb{R}^d), \quad \forall\, |\alpha| \geq 2.$$

Denote by $U(t) = e^{-itH_0}$ the linear flow map, and let $T > 0$.

(1) *For any admissible pair (p, q), there exists C_q such that*

$$\|U(t)\varphi\|_{L^p([0,T];L^q)} \leq C_q \|\varphi\|_{L^2}, \quad \forall\, \varphi \in L^2(\mathbb{R}^d).$$

(2) *Denote*

$$D(F)(t,x) = \int_0^t U(t-\tau) F(\tau, x) d\tau.$$

For all admissible pairs (p_1, q_1) and (p_2, q_2), there exists $C = C_{q_1,q_2}$ such that

$$\|D(F)\|_{L^{p_1}([0,t];L^{q_1})} \leq C \|F\|_{L^{p_2'}([0,t];L^{q_2'})}, \tag{3.1}$$

for all $F \in L^{p_2'}([0,T];L^{q_2'})$ and $0 \leq t \leq T$.

Remark 3.3. In general, the above constant C_q and C_{q_1,q_2} must be expected to depend on T. See, e.g., [4].

Duhamel's formulation for (1.1) takes the form

$$\psi_j(t) = U(t)\varphi_j - i\int_0^t U(t-\tau)\mathcal{C}_j(\psi_{-1},\psi_0,\psi_1)(\tau)d\tau$$
$$- i\int_0^t U(t-\tau)\mathcal{B}_j(\psi_{-1},\psi_0,\psi_1)(\tau)d\tau, \quad j=-1,0,1, \tag{3.2}$$

where the $\mathcal{C}_j$'s are $\mathbb{R}$-trilinear operators in their argument, and the $\mathcal{B}_j$ are $\mathbb{R}$-linear operators in their argument. Therefore, (3.2) is the system version of a cubic nonlinear equation, with a linear source term stemming from B. The classical theory for scalar nonlinear Schrödinger equations, as presented in, e.g., [4, 7], yields the following.

Proposition 3.4. *Suppose that B is bounded, $B \in L^\infty(\mathbb{R}^d)$, and that the external potential is smooth and at most quadratic*:

$$\partial^\alpha V_{\text{ext}} \in L^\infty(\mathbb{R}^d), \quad \forall\, |\alpha| \geq 2.$$

(1) *If $d=1$ and $\varphi_j \in L^2(\mathbb{R})$, then there exists T depending only on the $\|\varphi_j\|_{L^2}$'s, such that* (3.2) *has a unique solution*

$$\psi_j \in C([-T,T];L^2(\mathbb{R})) \cap L^8([-T,T];L^4(\mathbb{R})), \quad j=-1,0,1.$$

If in addition $B \in W^{1,\infty}(\mathbb{R})$ and $\varphi_j \in \widehat{\Sigma}$, then

$$\psi_j, \partial_x\psi_j, x\psi_j \in C([-T,T];L^2(\mathbb{R})) \cap L^8([-T,T];L^4(\mathbb{R})),$$
$$j=-1,0,1.$$

(2) *If $2 \leq d \leq 3$, suppose that $B \in W^{1,\infty}(\mathbb{R}^d)$. If $\varphi_j \in \widehat{\Sigma}$, then there exists T depending only on the $\|\varphi_j\|_{\widehat{\Sigma}}$'s, such that* (3.2) *has a unique solution*

$$\psi_j, \nabla\psi_j, x\psi_j \in C([-T,T];L^2(\mathbb{R}^d)) \cap L^{8/d}([-T,T];L^4(\mathbb{R}^d)),$$
$$j=-1,0,1.$$

(3) *In all the above cases, the conservation* (1.3) *holds, as well as* (1.4) *provided that $\varphi_j \in \widehat{\Sigma}$.*

Remark 3.5. If V_{ext} is at most linear in $x(\nabla V_{\text{ext}} \in L^\infty(\mathbb{R}^d))$, then the assumption $\varphi_j \in \widehat{\Sigma}$ can be relaxed to $\varphi_j \in H^1(\mathbb{R}^d)$, and the solution satisfies

$$\psi_j \in C([-T,T];H^1(\mathbb{R}^d)) \cap L^{8/d}([-T,T];W^{1,4}(\mathbb{R}^d)), \quad j=-1,0,1.$$

On the other hand, if V_{ext} is quadratic as in (1.2), working in $\widehat{\Sigma}$ (and not only in $H^1(\mathbb{R}^d)$) is necessary. See [5].

Corollary 3.6. *Suppose that B is bounded, $B \in L^\infty(\mathbb{R}^d)$, and that the external potential is smooth and at most quadratic.*

(1) *If $d=1$ and $\varphi_j \in L^2(\mathbb{R})$, then* (3.2) *has a unique global solution*

$$\psi_j \in C(\mathbb{R};L^2(\mathbb{R})) \cap L^8_{\text{loc}}(\mathbb{R};L^4(\mathbb{R})), \quad j=-1,0,1.$$

Commun. Contemp. Math. Downloaded from www.worldscientific.com by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.

If in addition $B \in W^{1,\infty}(\mathbb{R})$, *and* $\varphi_j \in \widehat{\Sigma}$, *then*

$$\psi_j, \partial_x \psi_j, x\psi_j \in C(\mathbb{R}; L^2(\mathbb{R})) \cap L^8_{\rm loc}(\mathbb{R}; L^4(\mathbb{R})), \quad j = -1, 0, 1.$$

(2) *If* $2 \leq d \leq 3$, *suppose that* $B \in W^{1,\infty}(\mathbb{R}^d)$. *If* $\varphi_j \in \widehat{\Sigma}$, *and* $\beta_s, \beta_n \geq 0$, *then*

$$\psi_j, \nabla \psi_j, x\psi_j \in C(\mathbb{R}; L^2(\mathbb{R}^d)) \cap L^{8/d}_{\rm loc}(\mathbb{R}; L^4(\mathbb{R}^d)), \quad j = -1, 0, 1.$$

Proof. Only the second point requires an explanation. From Proposition 3.4, we infer that (3.2) has a unique maximal solution. It is maximal in the future in the sense that either it is defined for all positive time, or there exists a finite T_+ such that

$$\|\psi_j(t)\|_{\widehat{\Sigma}} \xrightarrow[t \to T_+]{} \infty,$$

for at least one j. We then proceed as in [6], and introduce a modified energy functional

$$\begin{aligned}\mathcal{E}(\psi) = \int_{\mathbb{R}^d} \Bigg(& \sum_{j=-1}^{1} \left(\frac{1}{2} |\nabla \psi_j|^2 + \frac{|x|^2}{2} |\psi_j|^2 \right) + \frac{\beta_n}{2} \left(\sum_{j=-1}^{1} |\psi_j|^2 \right)^2 \\ & + \frac{\beta_s}{2} (|\psi_1|^2 - |\psi_{-1}|^2)^2 + \beta_s |\psi_0|^2 (|\psi_1|^2 + |\psi_{-1}|^2) \\ & + 2\beta_s \,\mathrm{Re}(\bar{\psi}_1 \psi_0^2 \bar{\psi}_{-1}) + 2\,\mathrm{Re}(B(\bar{\psi}_1 \psi_0 + \bar{\psi}_0 \psi_{-1})) \Bigg) dx.\end{aligned}$$

Since

$$2\,|\,\mathrm{Re}(\bar{\psi}_1 \psi_0^2 \bar{\psi}_{-1})| \leq |\psi_0|^2 (|\psi_1|^2 + |\psi_{-1}|^2)$$

and since the total mass is conserved, (1.3), in the case $\beta_s \geq 0$, we have

$$\begin{aligned}\mathcal{E}(\psi) \geq \int_{\mathbb{R}^d} \Bigg(& \sum_{j=-1}^{1} \left(\frac{1}{2} |\nabla \psi_j|^2 + \frac{|x|^2}{2} |\psi_j|^2 \right) + \frac{\beta_n}{2} \left(\sum_{j=-1}^{1} |\psi_j|^2 \right)^2 \\ & + 2\,\mathrm{Re}(B(\bar{\psi}_1 \psi_0 + \bar{\psi}_0 \psi_{-1})) \Bigg) dx.\end{aligned}$$

Since B is bounded, (1.3) yields

$$2 \left| \mathrm{Re} \int_{\mathbb{R}^d} (B(\bar{\psi}_1 \psi_0 + \bar{\psi}_0 \psi_{-1})) dx \right| \leq C,$$

for some uniform constant C. Therefore, in the case $\beta_n \geq 0$, there exists C such that

$$\mathcal{E}(\psi) \geq \frac{1}{2} \sum_{j=-1}^{1} \|\psi_j(t)\|^2_{\widehat{\Sigma}} - C. \tag{3.3}$$

Commun. Contemp. Math. Downloaded from www.worldscientific.com by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.

On the other hand, we compute, in view of (1.4):

$$\frac{d}{dt}\mathcal{E}(\psi(t)) = \frac{d}{dt}(\mathcal{E}(\psi(t)) - E(\psi(t)))$$
$$= \sum_{j=-1}^{1} \frac{d}{dt}\int_{\mathbb{R}^d}\left(\frac{|x|^2}{2} - V_{\rm ext}(x)\right)|\psi_j(t,x)|^2 dx.$$

For a general potential $\mathtt{V}(x)$, let

$$\mathcal{V}(t) := \sum_{j=-1}^{1}\int_{\mathbb{R}^d}\mathtt{V}(x)|\psi_j(t,x)|^2 dx.$$

We compute

$$\frac{d}{dt}\mathcal{V}(t) = 2\sum_{j=-1}^{1}\mathrm{Im}\int_{\mathbb{R}^d}\mathtt{V}(x)\bar{\psi}_j(t,x)i\partial_t\psi_j(t,x)dx.$$

In the same fashion as the total mass is conserved, algebraic cancellations yield

$$\frac{d}{dt}\mathcal{V}(t) = \sum_{j=-1}^{1}\mathrm{Im}\int_{\mathbb{R}^d}\bar{\psi}_j(t,x)\nabla\mathtt{V}(x)\cdot\nabla\psi_j(t,x)dx.$$

Since derivatives of order at least two of $V_{\rm ext}$ are bounded, there exists $C>0$ such that

$$|\nabla V_{\rm ext}(x)| \leq C\left(1+|x|\right), \quad \forall\, x\in\mathbb{R}^d.$$

We infer

$$\frac{d}{dt}\mathcal{E}(\psi(t)) \leq C\sum_{j=-1}^{1}\int_{\mathbb{R}^d}(1+|x|)|\bar{\psi}_j(t,x)\nabla\psi_j(t,x)|dx \leq C\sum_{j=-1}^{1}\|\psi_j(t)\|_{\Sigma}^2,$$

where the last estimate stems from Cauchy–Schwarz inequality. In view of (3.3), this rules out the existence of such a T_+, and the solution is global. □

3.1. *Virial*

In this section, we show that without the assumption $\beta_s, \beta_n \geq 0$, finite time blow-up may occur. Set

$$\mathcal{V}(t) := \sum_{j=-1}^{1}\int_{\mathbb{R}^d}|x|^2|\psi_j(t,x)|^2 dx. \tag{3.4}$$

From the above computations, we have

$$\dot{\mathcal{V}}(t) = 2\sum_{j=-1}^{1}\mathrm{Im}\int_{\mathbb{R}^d}\bar{\psi}_j(t,x)x\cdot\nabla\psi_j(t,x)dx.$$

Therefore,

$$\ddot{\mathcal{V}}(t) = 2\sum_{j=-1}^{1}\mathrm{Im}\int_{\mathbb{R}^d}(\partial_t\bar{\psi}_j(t,x)x\cdot\nabla\psi_j(t,x) + \bar{\psi}_j(t,x)x\cdot\nabla\partial_t\psi_j(t,x))dx.$$

Commun. Contemp. Math. Downloaded from www.worldscientific.com by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.

Integrating by parts, we have

$$\begin{aligned}
&\mathrm{Im}\int_{\mathbb{R}^d}\bar{\psi}_j(t,x)x\cdot\nabla\partial_t\psi_j(t,x)dx\\
&\quad=\sum_{k=1}^{d}\mathrm{Im}\int_{\mathbb{R}^d}\bar{\psi}_j(t,x)x_k\partial_k\partial_t\psi_j(t,x)dx\\
&\quad=-\sum_{k=1}^{d}\mathrm{Im}\int_{\mathbb{R}^d}\partial_t\psi_j(t,x)\partial_k(x_k\bar{\psi}_j(t,x))dx\\
&\quad=-\mathrm{Im}\int_{\mathbb{R}^d}\partial_t\psi_j(t,x)x\cdot\nabla\bar{\psi}_j(t,x)dx-d\,\mathrm{Im}\int_{\mathbb{R}^d}\bar{\psi}_j(t,x)\partial_t\psi_j(t,x)dx.
\end{aligned}$$

Summing over j, we infer

$$\ddot{\mathcal{V}}(t)=-2\sum_{j=-1}^{1}\mathrm{Im}\int_{\mathbb{R}^d}\partial_t\psi_j(t,x)(2x\cdot\nabla\bar{\psi}_j(t,x)+d\bar{\psi}_j(t,x))dx.$$

Lemma 3.7. *Let $\psi\in C(I;\widehat{\Sigma})^3$ be given by Proposition* 3.4, *for some time interval $I\ni 0$. The virial $\mathcal{V}$, defined by* (3.4), *satisfies*

$$\begin{aligned}
\ddot{\mathcal{V}}(t)&=2\sum_{j=-1}^{1}\|\nabla\psi_j\|_{L^2}^2-2\sum_{j=-1}^{1}\int|\psi_j|^2x\cdot\nabla V_{\mathrm{ext}}(x)dx+d\beta_n\int\left(\sum_{j=-1}^{1}|\psi_j|^2\right)^2dx\\
&\quad+d\beta_s\int((|\psi_{-1}|^2-|\psi_1|^2)^2+2|\psi_0|^2(|\psi_{-1}|^2+|\psi_1|^2)+4\,\mathrm{Re}(\bar{\psi}_{-1}\psi_0^2\bar{\psi}_1))dx\\
&\quad-4\,\mathrm{Re}\int x\cdot\nabla B(\psi_0\bar{\psi}_1+\psi_{-1}\bar{\psi}_0)dx,
\end{aligned}$$

which can also be written, in terms of the total (conserved) energy,

$$\begin{aligned}
\ddot{\mathcal{V}}(t)&=4E(\psi)-\sum_{j=-1}^{1}\int|\psi_j|^2(V_{\mathrm{ext}}+2x\cdot\nabla V_{\mathrm{ext}}(x))dx\\
&\quad+\beta_n(d-2)\int\left(\sum_{j=-1}^{1}|\psi_j|^2\right)^2dx+\beta_s(d-2)\int((|\psi_{-1}|^2-|\psi_1|^2)^2\\
&\quad+2|\psi_0|^2(|\psi_{-1}|^2+|\psi_1|^2)+4\,\mathrm{Re}(\bar{\psi}_{-1}\psi_0^2\bar{\psi}_1))dx\\
&\quad-2\,\mathrm{Re}\int(B+2x\cdot\nabla B)(\psi_0\bar{\psi}_1+\psi_{-1}\bar{\psi}_0)dx.
\end{aligned}$$

We readily infer that if $\beta_n,\beta_s<0$, and $d=2$ or 3, one may have $E(\psi)<0$ and, in the case where V_{ext} is harmonic, (1.2),

$$\ddot{\mathcal{V}}(t)\le 4E(\psi)-2\,\mathrm{Re}\int(B+2x\cdot\nabla B)(\psi_0\bar{\psi}_1+\psi_{-1}\bar{\psi}_0)dx.$$

Commun. Contemp. Math. Downloaded from www.worldscientific.com by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.

With $B \in W^{1,\infty}$,

$$\begin{aligned}\ddot{\mathcal{V}}(t) &\leq 4E(\psi) + \int (\|B\|_{L^\infty} + |x|\|\nabla B\|_{L^\infty})\|\psi_0\bar{\psi}_1 + \psi_{-1}\bar{\psi}_0\|dx \\ &\leq 4E(\psi) + (\|B\|_{L^\infty}\|\psi_0\|_{L^2} + \|\nabla B\|_{L^\infty}\|x\psi_0\|_{L^2})(\|\psi_1\|_{L^2} + \|\psi_{-1}\|_{L^2}) \\ &\leq 4E(\psi) + (\|B\|_{L^\infty}\|\psi_0\|_{L^2} + \|\nabla B\|_{L^\infty}\sqrt{\mathcal{V}(t)})(|\psi_1\|_{L^2} + \|\psi_{-1}\|_{L^2}).\end{aligned}$$

From this relation, it is clear that this inequality may lead to the cancellation of $\mathcal{V}$, which is not possible in the case of a solution with $\widehat{\Sigma}$ regularity, for which the total mass is conserved. Since the main goal of this paper is to study the orbital stability of standing waves in the case $\beta_s, \beta_n \geq 0$, we shall not pursue on the possibility of finite time blow-up.

4. Orbital Stability

As in the previous sections, we suppose that $\beta_s, \beta_n > 0$.

Theorem 4.1. *Let $M, N > 0$.*

(A) *If $B \equiv 0$*

(a) *$I_{M,N} = \widehat{I}_{M,N}, Z_{M,N} \neq \emptyset$ and $Z_{M,N}$ is stable.*

(b) *For any $z = (z_{-1}, z_0, z_1) \in Z_{M,N}$, we have $|z| = (|z_{-1}|, |z_0|, |z_1|) \in W_{M,N}$ and*

$$Z_{M,N} = \{e^{i\theta_1}w_{-1}, e^{i\theta_2}w_0, e^{i\theta_3}w_1; w \in W_{M,N}, (\theta_1, \theta_2, \theta_3) \in \mathbb{R}^3\}$$

(w_{-1}, w_0, w_1) is a solution of (1.10) and if

$$w_{-1} = |w_{-1}|e^{i\theta_{-1}}, w_0 = |w_0|e^{i\theta_0} \quad \text{and} \quad w_1 = |w_1|e^{i\theta_1}$$

then

$$2\theta_0 - \theta_1 - \theta_{-1} = \pi \mod 2\pi. \tag{4.1}$$

(B) *If $B \not\equiv 0$*

(a) *$Z_{M,N}$ is stable.*

(b) *$\widehat{I}_{M,N} > I_{M,N}$.*

Proof. (A) If $B \equiv 0$:

(a): As suggested in [8], to prove the orbital stability of standing waves of (1.1), it suffices to prove that $Z_{M,N} \neq \emptyset$ and any minimizing sequence,

$$z_n = (z_{n,-1}, z_{n,0}, z_{n,1}) \in \widehat{\Sigma}^3 \text{ such that } z_n \in \widehat{C}_{M,N} \text{ and } \widehat{E}(z_n) \to \widehat{I}_{M,N} \tag{4.2}$$

is relatively compact in $\widehat{\Sigma}^3$. Consider a sequence such that $z_n \in \widehat{C}_{M,N}$ and $\widehat{E}(z_n) \to \widehat{I}_{M,N}$. The first goal is to show that $\{z_n\}$ has a subsequence which is convergent in $\widehat{\Sigma}^3$. By Lemma 2.3, we see that z_n is bounded in $\widehat{\Sigma}^3$. Therefore passing to a subsequence, we can suppose that:

$$u_{n,j} \rightharpoonup u_j \text{ and } v_{n,j} \rightharpoonup v_j \text{ in } \Sigma, -1 \leq j \leq 1. \tag{4.3}$$

Commun. Contemp. Math. Downloaded from www.worldscientific.com by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.

Now let $\rho_{n,j} = |z_{n,j}| = (u_{n,j}^2 + v_{n,j}^2)^{1/2}, z_{n,j} = \rho_{n,j} e^{i\theta_{n,j}}, -1 \leq j \leq 1$. It follows that $\rho_{n,j} \in \Sigma$ and that for all $n \in \mathbb{N}$ and $-1 \leq j \leq 1, 1 \leq k \leq d$,

$$\partial_k \rho_{n,j}(x) = \begin{cases} \dfrac{u_{n,j}(x)\partial_k u_{n,j}(x) + v_{n,j}(x)\partial_k v_{n,j}(x)}{(u_{n,j}(x)^2 + v_{n,j}(x)^2)^{1/2}} & \text{if } u_{n,j}(x)^2 + v_{n,i}(x)^2 > 0, \\ 0 & \text{otherwise.} \end{cases}$$

Therefore

$$\widehat{E}(z_n) - E(\rho_n) = \frac{1}{2}(\|\nabla z_n\|_2^2 - \|\nabla |z_n|\|_2^2) + 2\beta_s \int_{\mathbb{R}^d} |z_{n,1}||z_{n,0}|^2 |z_{n,-1}| \times (\cos(2\theta_{n,0} - \theta_{n,-1} - \theta_{n,1}) + 1). \tag{4.4}$$

The triangle inequality

$$||z_n(t, x+h)| - |z_n(t,x)|| \leq |z_n(t, x+h) - z_n(t,x)|$$

implies

$$\|\nabla z_n\|_2^2 - \|\nabla |z_n|\|_2^2 \geq 0. \tag{4.5}$$

Therefore $\widehat{E}(z_n) - E(\rho_n) \geq 0$, hence $\widehat{I}_{M,N} = \lim_{n\to\infty} \widehat{E}(z_n) \geq \limsup E(\rho_n)$. Taking into account that

$$\|z_{n,j}\|_2^2 = \|\rho_{n,j}\|_2^2 =: c_{n,j}^2, \quad \text{for all } -1 \leq j \leq 1$$

and that $\widehat{I}_{M,N} \leq I_{M,N}$, it follows by using Lemma 2.3 that

$$\liminf E(\rho_n) \geq \liminf I_{M_n,N_n} \geq I_{M,N} \geq \widehat{I}_{M,N},$$
$$M_n = c_{n,1}^2 + c_{n,0}^2 + c_{n,-1}^2 \quad \text{and} \quad N_n = c_{n,1}^2 - c_{n,-1}^2.$$

Therefore

$$\lim_{n\to\infty} E(\rho_n) = \lim_{n\to\infty} \widehat{E}(z_n) = I_{M,N} = \widehat{I}_{M,N}. \tag{4.6}$$

Now on the other hand (4.5) implies that for any $-1 \leq j \leq 1$ we have

$$\lim_{n\to\infty} \int_{\mathbb{R}^d} |\nabla u_{n,j}|^2 + |\nabla v_{n,j}|^2 - |\nabla (u_{n,j}^2 + v_{n,j}^2)^{1/2}|^2 = 0. \tag{4.7}$$

Thus:

$$\lim_{n\to\infty} \int_{\mathbb{R}^d} |\nabla u_{n,j}|^2 + |\nabla v_{n,j}|^2 = \lim_{n\to\infty} \int_{\mathbb{R}^d} |\nabla (u_{n,j}^2 + v_{n,j}^2)^{1/2}|^2, \tag{4.8}$$

which is equivalent to saying that

$$\lim_{n\to\infty} \|\nabla z_n\|_2^2 = \lim_{n\to\infty} \|\nabla |z_n|\|_2^2. \tag{4.9}$$

Commun. Contemp. Math. Downloaded from www.worldscientific.com by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.

Using (4.5)–(4.6) and Lemma 2.3, $\rho_n = (\rho_{n,-1}, \rho_{n,0}, \rho_{n,1})$ is relatively compact in Σ^3. Thus there exist $\rho_{-1}, \rho_0, \rho_1 \in \Sigma$ such that

$$\begin{cases} (u_{n,j}^2 + v_{n,j}^2)^{1/2} \text{ converges to } \rho_j \text{ in } \Sigma, \quad -1 \le j \le 1, \\ |\rho_i|_2 = c_j, \quad \text{with } \sum_{j=-1}^{1} c_j^2 = M \text{ and } c_1^2 - c_{-1}^2 = N, \\ E(\rho_{-1}, \rho_0, \rho_1) = I_{M,N}. \end{cases} \tag{4.10}$$

Closely following [10], we will prove that $\rho_j = |z_j| = (u_j^2 + v_j^2)^{1/2}$. By (4.3), we know that $u_{n,j} \to u_j$ and $v_{n,j} \to v_j$ in $L^2(B(0,R))$ we can easily deduce that

$$[(u_{n,j}^2 + v_{n,j}^2)^{1/2} - (u_j^2 + v_j^2)^{1/2}]^2 \le |u_{n,j} - u_j|^2 + |v_{n,j} - v_j|^2,$$

thus $(u_{n,j}^2 + v_{n,j}^2)^{1/2} \to \rho_j$ in $L^2(B(0,R))$, this certainly implies that $|z_j| = \rho_j$ for all $-1 \le j \le 1$. On the other hand

$$\|z_{n,j}\|_2 = \||z_{n,j}|\|_2 \to c_j = \|z_j\|_2 = \||z_j|\|_2.$$

The proof of the first part of Theorem 4.1 is now complete if we show that

$$\lim_{n\to\infty} \|\nabla z_{n,j}\|_2^2 = \|\nabla z_j\|_2^2, \quad -1 \le j \le 1.$$

From (4.7), we have

$$\lim_{n\to\infty} \|\nabla z_{n,j}\|_2 = \lim_{n\to\infty} \||\nabla z_{n,j}|\|_2$$

and

$$\lim_{n\to\infty} |\nabla |z_{n,j}||_2 = |\nabla |z_j||_2.$$

Thus by the lower semi-continuity of $|\cdot|_2$, we have

$$\|\nabla z_i\|_2^2 \le \lim \|\nabla z_i\|_2 = \lim |\nabla |z_{n,j}||_2^2 = |\nabla |z_j||_2^2. \tag{4.11}$$

Finally, replacing $z_{n,j}$ by z_j in (4.5), we see that

$$\|\nabla z_j\|_2^2 \ge \||\nabla |z_j|\|_2^2 \quad \text{for all} -1 \le j \le 1.$$

Now using (4.3), we know that $z_{n,j} \to z_j$ in $\widehat{\Sigma}$.

(b): Let $z = (z_1, z_0, z_1) \in Z_{M,N}$ with

$$z_j = |z_j| e^{i\theta_j} = (u_j, v_j), \quad -1 \le j \le 1.$$

By (4.5), we know that for all $-1 \le j \le 1, 1 \le k \le d$,

$$\begin{cases} \displaystyle\int_{\mathbb{R}^d} \frac{(u_j \partial_k v_j - v_j \partial_k u_j)^2}{u_j^2 + v_j^2} dx = 0, \\ 2\theta_0 - \theta_1 - \theta_{-1} = \pi \mod 2\pi. \end{cases} \tag{4.12}$$

Commun. Contemp. Math. Downloaded from www.worldscientific.com by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.

On the other hand $\widehat{E}(z) = \widehat{I}_{M,N}$, which implies that there exists a Lagrange multiplier $\alpha \in \mathbb{C}$ such that

$$\widehat{E}'(z)\xi = \frac{\alpha}{2} \sum_{i=-1}^{1} z_i \bar{\xi}_i + \xi_i \bar{z}_i \quad \text{for all } \xi \in \mathbb{C}^3.$$

By elementary regularity theory and maximum principle, we can prove that u_j and $v_j \in C^1(\mathbb{R}^d) \cap \Sigma$ and $\rho > 0$ (see [14, Sec. 4]).

Set $\Omega_j = \{x \in \mathbb{R}^2 : u_j(x) = 0\}$. Then Ω_j is closed since u_j is continuous. Let us prove that it is also open. Let $x_0 \in \Omega_j$. Using the fact that $v_j(x_0) > 0$, we can find a ball B centered in x_0 such that $v_j(x) \neq 0$ for any $x \in B$. Consequently for $-1 \leq j \leq 1$, and $1 \leq k \leq d$,

$$\forall\, x \in B, \quad \left(\frac{(u_j \partial_k v_j - v_j \partial_k v_j)^2}{u_j^2 + v_j^2} \right) = \left(\left(\partial_k \left(\frac{u_j}{v_j} \right) \right)^2 \frac{v_j^4}{u_j^2 + v_j^2} \right).$$

This implies that

$$\int_B \left| \nabla \left(\frac{u_j}{v_j} \right) \right|^2 \frac{v_j^4}{u_j^2 + v_j^2} = 0. \tag{4.13}$$

Hence $\nabla(\frac{u_j}{v_j}) = 0$ on B, and there exists c such that $\frac{u_j}{v_j} = c$ on B. Since $x_0 \in B$, we infer $c \equiv 0$. Therefore Ω_j is also an open set of $\mathbb{R}^d$. Hence we have proved that for $-1 \leq j \leq 1$, we have these two alternatives

$$A1 : u_j \equiv 0 \text{ or } u_j > 0 \quad \text{on } \mathbb{R}^d,$$

$$A2 : v_j \equiv 0 \text{ or } v_j > 0 \quad \text{on } \mathbb{R}^d.$$

Now let $z_j = e^{i\sigma_j} w_j, \sigma \in \mathbb{R}$, $w_j \in W_{M,N}$:

$$\sum_{j=-1}^{1} |z_j|_2^2 = N, \quad \text{and} \quad |z_1|_2^2 - |z_{-1}|_2^2 = M.$$

Obviously

$$\{e^{i\sigma_j} w_j; \sigma_j \in \mathbb{R}, w \in W_{M,N}\} \subset Z_{M,N}.$$

Conversely for $z_j = (u_j, v_j)$ such that $(z_{-1}, z_0, z_1) \in Z_{M,N}$, $\widehat{E}(z_1, z_0, z_1) = \widehat{E}(z) = \widehat{I}_{M,N}$, with $(w_{-1}, w_0, w_1) \in W_{M,N}$. Now we have six possible alternatives. We discuss one them in details.

Suppose that v_1, v_0 and $v_{-1} \neq 0$ for all $x \in \mathbb{R}^d$. In this case, it follows that $\nabla(\frac{u_j}{v_j}) = 0$ on $\mathbb{R}^d$. Thus we can find three constants K_{-1}, K_0, K_1 such that

$$u_j = K_j v_j, \quad -1 \leq j \leq 1.$$

Therefore $w_j = (K_j + i)v_j$. Let $\theta_j \in \mathbb{R}$ such that $K_j + i = |K_j + i| e^{i\theta_j}$ and let $\eta_j = 0$ if $v_j > 0$ and $\eta_j = \pi$ if $v_j < 0$ on $\mathbb{R}^d$. Setting, $\sigma_j = \theta_j + \eta_j$, we infer

$$z_j = (K_j + i)v_j = |K_j + i| e^{i\theta_j} |v_j| e^{i\eta_j} = w_j e^{i\sigma_j}.$$

Commun. Contemp. Math. Downloaded from www.worldscientific.com by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.

(B) If $B \not\equiv 0$:

(a): It is sufficient to prove that any minimizing sequence is relatively compact in $\widehat{\Sigma}^3$. The proof goes exactly the same way as in Lemma 2.3.

(b): We will argue by contradiction. Suppose that $\widehat{I}_{M,N} = I_{M,N}$, let (z_n) denote a minimizing sequence of the constrained variational problem (1.11). All the steps of the proof of (A)(a) remain true until (4.4) which becomes now:

$$\begin{aligned}\widehat{E}(z_n) - E(\rho_n) = \frac{1}{2}(\|\nabla z_n\|_2^2 - \|\nabla |z_n|\|_2^2) \\ + 2\beta_s \int_{\mathbb{R}^d} |z_{n,1}||z_{n,0}|^2 |z_{n,-1}|(\cos(2\theta_{n,0} - \theta_{n,-1} - \theta_{n,1}) + 1). \\ + 2\int_{\mathbb{R}^d} |z_{n,1}||z_{n,0}|(\cos(\theta_{n,0} - \theta_{n,1}) + 1) \\ + 2\int_{\mathbb{R}^d} |z_{n,-1}||z_{n,0}|(\cos(\theta_{n,-1} - \theta_{n,0}) + 1); \end{aligned} \tag{4.14}$$

and (4.12) is now:

$$\begin{cases} \displaystyle\int_{\mathbb{R}^d} \frac{(u_j\partial_k v_j - v_j\partial_k u_j)^2}{u_j^2 + v_j^2} dx = 0, \\ 2\theta_0 - \theta_1 - \theta_{-1} = \pi \mod 2\pi, \\ \theta_0 - \theta_1 = \pi \mod 2\pi, \\ \theta_0 + \theta_{-1} = \pi \mod 2\pi, \end{cases} \tag{4.15}$$

which implies that $2\theta_0 - \theta_1 - \theta_{-1} = \pi \mod 2\pi$ and $2\theta_0 - \theta_1 - \theta_{-1} = 0 \mod 2\pi$. Thus, we get our contradiction.

Remark 4.2. We still have no clue about the characterization of the orbit of standing waves. We believe that a new approach must be built up to solve this challenging problem. This new phenomena ($\widehat{I}_{M,N} > I_{M,N}$) appears in many other Bose–Einstein systems. □

Acknowledgment

H. Hajaiej would like to thank the National Plan of Science and Technology, NPST, for its support through the research project 11-MAT1716-02.

References

[1] W. Bao, I.-L. Chern and Y. Zhang, Efficient numerical methods for computing ground states of spin-1 Bose–Einstein condensates based on their characterizations, *J. Comput. Phys.* **253** (2013) 189–208.

[2] A. Burchard and H. Hajaiej, Rearrangement inequalities for functionals with monotone integrands, *J. Funct. Anal.* **233**(2) (2006) 561–582.

Commun. Contemp. Math. Downloaded from www.worldscientific.com by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.

[3] D. Cao, I.-L. Chern and J.-C. Wei, On ground state of spinor Bose–Einstein condensates, *NoDEA Nonlinear Differential Equations Appl.* **18**(4) (2011) 427–445.
[4] R. Carles, Linear vs. nonlinear effects for nonlinear Schrödinger equations with potential, *Commun. Contemp. Math.* **7**(4) (2005) 483–508.
[5] ———, On the Cauchy problem in Sobolev spaces for nonlinear Schrödinger equations with potential, *Portugal. Math.* (*N.S.*) **65**(2) (2008) 191–209.
[6] R. Carles and J. Drumond Silva, Large time behavior in nonlinear Schrödinger equation with time dependent potential, *Commun. Math. Sci.* **13**(2) (2015) 443–460.
[7] T. Cazenave, *Semilinear Schrödinger Equations*, Courant Lecture Notes in Mathematics, Vol. 10 (University Courant Institute of Mathematical Sciences, New York, 2003).
[8] T. Cazenave and P.-L. Lions, Orbital stability of standing waves for some nonlinear Schrödinger equations, *Comm. Math. Phys.* **85**(4) (1982) 549–561.
[9] U. Ernst, A. Marte, F. Schreck, J. Schuster and G. Rempe, Bose–Einstein condensation in a pure Ioffe–Pritchard field configuration, *Europhys. Lett.* **41** (1998) 1–6.
[10] H. Hajaiej and C. A. Stuart, On the variational approach to the stability of standing waves for the nonlinear Schrödinger equation, *Adv. Nonlinear Stud.* **4**(4) (2004) 469–501.
[11] T. Isoshima, M. Nakahara, T. Ohmi and K. Machida, Creation of a persistent current and vortex in a Bose–Einstein condensate of alkali-metal atoms, *Phys. Rev. A* **61**(6) (2000) 063610.
[12] O. Kavian and F. Weissler, Self-similar solutions of the pseudo-conformally invariant nonlinear Schrödinger equation, *Michigan Math. J.* **41**(1) (1994) 151–173.
[13] A. E. Leanhardt, Y. Shin, D. Kielpinski, D. E. Pritchard and W. Ketterle, Coreless vortex formation in a spinor Bose–Einstein condensate, *Phys. Rev. Lett.* **90** (2003) 140403.
[14] L. Lin and I.-L. Chern, Bifurcation between 2-component and 3-component ground states of spin-1 Bose-Einstein condensates in uniform magnetic fields, preprint (2013); http://arXiv.org/abs/1302.0279.

Commun. Contemp. Math. Downloaded from www.worldscientific.com
by UNIVERSITY OF SUSSEX on 06/22/16. For personal use only.